



BRIEF REPORT

An ice-binding protein from an Arctic grass, *Leymus mollis*

[version 1; peer review: 1 approved]

Todd L. Sformo ^{1,2}, James A. Raymond ³

¹Department of Wildlife Management, North Slope Borough, Utqiagvik, Alaska, 99723, USA

²Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, Alaska, 99775, USA

³School of Life Sciences, University of Nevada Las Vegas, Las Vegas, Nevada, 89154, USA

v1 First published: 26 Jun 2020, 9:648
<https://doi.org/10.12688/f1000research.24328.1>

Latest published: 26 Jun 2020, 9:648
<https://doi.org/10.12688/f1000research.24328.1>

Abstract

Several cold-hardy grasses have been shown to have ice-binding proteins (IBPs) that protect against freeze-thaw injury. Here, we looked for IBP activity in an Alaskan coastal grass that had not previously been examined, *Leymus mollis* (Pooidae). Rhizome tissue had strong ice-structuring and ice recrystallization inhibiting (IRI) activities, indicating the probable presence of IBPs. The gene sequence of an IBP was obtained. The sequence encoded a 118-amino acid IRI domain that contained eight repeats. A 3D structure of the IRI domain was predicted from the structure of the IRI domain of the perennial ryegrass *Lolium perenne*. The predicted structure appeared to have the same eight beta-roll coils found in the *L. perenne* IBP.

Keywords

Leymus mollis, ice-binding protein, Arctic, Alaska, Pooidae, dune grass, Beaufort Sea

Open Peer Review

Reviewer Status

Invited Reviewers

1

version 1

26 Jun 2020

report

1 Peter L. Davies , Queen's University,
Kingston, Canada

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: James A. Raymond (raymond@unlv.nevada.edu)

Author roles: Sformo TL: Conceptualization, Funding Acquisition, Investigation, Resources, Validation, Writing – Review & Editing; Raymond JA: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Resources, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.




Grant information: The author(s) declared that no grants were involved in supporting this work.


Copyright: © 2020 Sformo TL and Raymond JA. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Sformo TL and Raymond JA. An ice-binding protein from an Arctic grass, *Leymus mollis* [version 1; peer review: 1 approved] F1000Research 2020, 9:648 <https://doi.org/10.12688/f1000research.24328.1>

First published: 26 Jun 2020, 9:648 <https://doi.org/10.12688/f1000research.24328.1>

Introduction

Within the grass family (Poaceae), the subfamily Pooideae includes many cold-adapted grasses including wheat, barley and forage grasses. These grasses have developed a large family of ice-binding proteins (IBPs) that protect the plants from freezing damage (Sandve *et al.*, 2008). The IBPs, which have negligible effect on the freezing point, are characterized by a C-terminal domain consisting of several repeating units that is associated with strong ice recrystallization inhibition (IRI) activity (Sandve *et al.*, 2008). This region is thus called the IRI domain. Ice recrystallization, in which larger ice grains grow at the expense of smaller grains, occurs mostly at warmer sub-zero temperatures and is thought to cause damage to cell walls. Two of the most studied of these IBPs are from the grasses *Lolium perenne* (Sidebottom *et al.*, 2000) and *Deschampsia antarctica* (John *et al.*, 2009). Their amino acid sequences are very similar in both the N-terminal and IRI domains. The 118-a.a. IRI domain of the *L. perenne* IBP has been crystallized and its structure determined by X-ray crystallography (Middleton *et al.*, 2012). The IRI domain has a beta-roll fold with eight similar coils, one side of which is the predicted ice-binding site. The spacing of the coils is very close to the repeat distance along the a-axis of ice. Interestingly, heterologous expression of the *Lolium* IRI domain in tomato was also shown to increase chilling (4°C) tolerance (Balamurugan *et al.*, 2018), although the mechanism remains unclear. Here, we describe a related IBP from a pooid grass from Utqiagvik (Barrow), Alaska, *Leymus mollis*. This site, at 71.3° N, is the highest latitude at which IBPs have been examined in a grass  is 9° higher in latitude than the site where *D. antarctica*  was sampled in the southern hemisphere. 

L. mollis, also known as American dune grass, is found in coastal habitats, especially sand dunes across North America and Greenland. It is typically subjected to many stresses, such as low nutrient levels, ~~salty sand~~, salt spray, little freshwater, inundation during storms, wind abrasion, and ice storms (Gagné & Houle, 2002).  *L. mollis*'s IBP sequence is very similar to the *Lolium* and *Deschampsia* sequences, but closest to wheat (*Triticum*) IBPs. We also show that an extract of the grass has strong ice-shaping and recrystallization inhibition properties and predict its structure based on the structure of *L. perenne* IBP.

Methods

Grass

A grass sample was collected from a gravel beach at Utqiagvik (Barrow), Alaska (71.3°N) on 6 October 2019, and stored at -80°C. The grass was shipped frozen to University of Nevada Las Vegas for analysis, but accidentally rose to ~10°C for about one day during shipment.

Activity measurements

To obtain a sample for measuring ice structuring and IRI activities, the brown outer layers of a rhizome from just below the soil surface were peeled off, revealing green tissue

underneath. About 70 mg of the green tissue was ground in 1 ml water in a mortar and pestle. The suspension was centrifuged at 14,000 rpm for 5 minutes to yield a slightly cloudy supernatant. Stem tissue from lawn grass (*Festuca* sp.) at the University of Nevada Las Vegas (52 mg) was similarly homogenized and used as a control. Ice-structuring activity was observed by examining the growth of an ice seed crystal with well-defined ice c- and a-axes submerged in the grass extract supernatant, as described previously (Raymond & Fritsen, 2001). Briefly, the sample was placed in a rectangular tube and the tube was submerged in a controlled temperature bath with front and rear windows. The growth of the crystal was observed at a temperature slightly below the freezing point (~-0.2°C) with a horizontally mounted dissecting microscope. Ice-structuring activity was defined as the appearance of sharply defined facets on the crystal surface. IRI activity was observed as described previously (Raymond & Fritsen, 2001). Briefly, 3 µl drops of supernatant were placed on a slide cover glass. A liquid nitrogen-cooled slide glued to a handle was pressed against the drops to form highly polycrystalline ice between the slide and cover glass. The samples were stored at -3°C in hexane and changes in recrystallization were monitored in the temperature bath described above over 24 hours. Photographs were taken through crossed polarizers.

Sequencing

Green tissue was obtained and homogenized as described above. DNA was extracted with a NucleoSpin Plant II kit (Machery Nagel; catalog number 740770) according to the manufacturer's instructions. To make IBP primers, the nucleotide sequences of IBPs from *L. perenne*, *D. antarctica* and *Triticum aestivum* (GenBank accession numbers EU680848, FJ663044 and KU204387, respectively) were aligned. Regions with high identities at the 5' and 3' ends were selected for primer sequences to obtain an amplicon that covered as much of the gene as possible. Primers for 18S ribosomal RNA were selected from conserved regions in the 18S sequences from *Dupontia fisheri*, *L. perenne* and *D. antarctica* (GenBank accession numbers KP794861, KJ598999 and MH628292, respectively) that flanked a region of high variability. At that time, we suspected the Utqiagvik grass sample was *D. fisheri* and didn't use the *L. mollis* 18S sequence. PCR was carried out with an Eppendorf Mastercycler Personal thermal cycler, with 3 minutes initial denaturation at 95°C, followed by 35 cycles of 30 seconds denaturation at 95°C, 30 seconds annealing at 59°C and 40 seconds extension at 72°C, followed by 3 minutes final extension at 72°C using Promega GoTaq polymerase (catalog number M300B). PCR products were electrophoresed on a 2% agarose gel, stained with ethidium bromide and observed and photographed with a UVP transilluminator. IBP and 18S bands of the expected sizes were obtained (see *Underlying data* (Raymond, 2020c)), cleaned up with a Nucleospin gel and PCR clean-up kit (Machery Nagel, catalog number 740609) and sequenced in both directions at the UNLV Genomics Core with an Applied Biosystems 3130 sequencer. The primers that were used for sequencing are shown in Table 1.

Table 1. Primers used in this study.

Primer pair	Fwd	Rev	Size (bp)
IBP	5'-TGCCACCCCGATGACCTCCG-3'	5'-TTAACCTCCTGTCACGACTTTGTTGCTCCC-3'	798
18S	5'-GGAAGGATCATTGTCGTGACCCTGACC-3'	5'-CTGGGGTCGCGGTCGAAGCGTC-3'	623

IBP, ice-binding protein.

Structure prediction

A 3D model of the IRI domain of the *Leymus* IBP was predicted with SWISS-MODEL (Waterhouse *et al.*, 2018) using the eight-coil structure of the *Lolium perenne* IRI domain (Protein Data Bank accession no. 3ULT) (Middleton *et al.*, 2012) as template. SWISS MODEL proposed three structures of the *Leymus* IRI domain based on the template. We selected the one with the highest sequence identity (76.5%), which also was the only one that had eight orderly coils like those in the *L. perenne* template. The free energy of this structure was then minimized (from -42.56 to -52.58 MJ mol⁻¹) with the YASARA Energy Minimization Server (Krieger *et al.*, 2009) and displayed with YASARA View. Stereoviews were obtained by rotating the molecule around its vertical axis by 3°. The average distance between coils in the IRI domain was measured with the YASARA distance function as the distance between the alpha carbons of Ser15 and Ser116 divided by seven.

Results and discussion

Species identification

The obtained 18S rRNA sequence (GenBank accession no. MT506010) most closely matched the sequence of *Leymus mollis* (GenBank accession no. EF581964) (98.3% identity). Among the Pooidae, *L. mollis* is a member of the Triticodae, which includes wheat (*Triticum*) and barley (*Hordeum*), while *Lolium* and *Deschampsia* are members of the Pooidae. The grass was confirmed as *L. mollis* from photos of the remnant spikes and leaves by Matthew Carlson (University of Alaska Anchorage) and Carolyn Parker, University of Alaska Fairbanks (personal communications to T.S.).

Ice-structuring and recrystallization inhibition activities

An extract from green rhizome tissue strongly affected the growth of an ice seed crystal, causing it to develop sharp facets (Figure 1A). The facets are an indication of the presence of ice-binding proteins. Extract from lawn grass stem tissue (control) showed no such activity (Figure 1B). The *Leymus* extract also had strong IRI activity. A polycrystalline ice crystal recrystallized significantly after 21 hours at -3°C, while the supernatant of the grass extract showed no recrystallization (Figure 2).

Ice-binding protein

Primers based on the IBPs of other pooid grasses succeeded in amplifying a sequence that encoded the C-terminal part of the IBP gene of *L. mollis*, including the entire IRI domain. The




Figure 1. Ice-structuring abilities of grass extracts. (A) *Leymus mollis* rhizome tissue. (B) Lawn grass (*Festuca* sp.) stem tissue (control). Ice seed crystals were placed in the extracts at slightly below the freezing point. Scale bars, 1 mm.




Figure 2. Comparison of ice recrystallization in 3 µl drops of water (ICE) and *L. mollis* extract (L.m.). Scale bar, 1 mm.

sequence (GenBank accession no. MT506011) encoded 260 a.a., which corresponds to all but the first 25 a.a. of the *L. perenne* IBP sequence. The sequence included the stop codon and contained no introns. Although the sequence was close to

the sequence of *L. perenne*, it most closely matched an IBP-like protein from *Triticum aestivum* (QBE94480) (86% identity, 91% similarity), in agreement with *L. mollis*'s classification as a member of the Triticoideae. When only the IRI domains were compared, the *Leymus* IRI domain was also closer to the *Triticum* IRI domain than the *Lolium* IRI domain. The ice-binding activities of *Triticum* spp. IBPs have not yet been reported.

The IRI domain of *L. mollis*, like that of *L. perenne*, has eight repeats, each consisting of 14 or 15 residues (Figure 3A). The consensus sequences of the two IBPs are virtually the same (Figure 3A).

Model of structure

The predicted structure of the *L. mollis* IBP IRI domain, obtained from using the *L. perenne* IBP IRI domain structure as template, resembles the *Lolium* IRI domain in several respects: it has eight repeats corresponding to eight coils in a beta-roll

fold (Figure 3B); a flat side (the a side) that is populated with two rows that are rich in Thr and Ser residues (Figure 3B); an irregular side (the b side) that has a bulge in the first three coils (Figure 3C); and an interior that is dominated by Asn/His ladders (red residues in Figure 3A). The a side has been identified as the ice-binding site of *L. perenne* IBP (Middleton *et al.* 2012). In *Lolium*, the average spacing between the alpha carbons of the row of amino acids on the ice-binding site is 4.5 Å, which is very similar to the spacing along the a-axis of ice (4.51 Å). In *Leymus*, the average spacing was calculated as 4.77 Å.

In summary, we describe an ice-binding protein from the grass *Leymus mollis* with ice-structuring and ice recrystallization inhibition (IRI) activities. The collection site of the grass, a gravelly beach on the Chukchi Sea, is an extreme habitat subject to numerous environmental stresses. *L. mollis* is also the highest latitude pooid grass so far examined for IBPs. Its IBP is similar in sequence to IBPs from other grasses and the predicted




Figure 3. Structure of the IRI domain of *Leymus mollis* IBP. (A) Comparison of the *Leymus* repeats (bottom) with those in the IRI domain of *Lolium perenne* (top). The *Lolium* data and the color scheme have been reproduced with permission from Middleton *et al.* (2012). Both domains have eight repeats. Consensus sequences are shown on top. Numbers indicate a.a. residue numbers. In the *Lolium* sequence, the gray background indicates the ice-binding site called the a side and the yellow background indicates a bulge on the b side. Similar features are found in the *Leymus* sequence. (B, C) Stereoviews predicted by SWISS MODEL using the IRI domain of *L. perenne* as template. SWISS MODEL was able to model the *Leymus* structure from Pro11 to Gly124, which corresponds to Asp1 to Ala118 in *L. perenne*. (B) View through the center of the coils, in which the ice-binding site (the a side) is on top. Amino acid side chains are shown only for the a and b sides. Color code of the ribbon: red, beta strand; cyan, coil. Color code of amino acid side chain: Cyan, C; blue, N; red, O; gray, H. (C) View of the b side of a space-filling model. A bulge is created by two residues on each of the first three coils. The first of the two residues in each coil is labeled. Colors are the same as those for the side chains in A.

3D structure of its IRI domain is very similar to that of the ryegrass *Lolium perenne*.

Data availability

Underlying data

Leymus mollis 18S ribosomal sequence on GenBank, Accession number [MT506010](#)

Leymus mollis IBP sequence on GenBank, Accession number [MT506011](#)

Figshare: Ice structuring by ice-binding protein of *Leymus mollis* DSCN5945.JPG. <https://doi.org/10.6084/m9.figshare.12401966.v1> (Raymond, 2020a)

Figshare: *Leymus mollis* ice recrystallization inhibition. <https://doi.org/10.6084/m9.figshare.12401954.v1> (Raymond, 2020b)

Figshare: PCR amplification of IBP and 18S genes of *Leymus mollis*. <https://doi.org/10.6084/m9.figshare.12401957.v1> (Raymond, 2020c)

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Acknowledgements

JR thanks the School of Life Sciences, University of Nevada Las Vegas, for providing facilities for carrying out this study.

References

Balamurugan S, Ann JS, Varghese IP, *et al.*: **Heterologous expression of *Lolium perenne* antifreeze protein confers chilling tolerance in tomato.** *J Integrative Agric.* 2018; **17**(5): 1128–1136.

[Publisher Full Text](#)

Gagné JM, Houle G: **Factors responsible for *Honckenya peploides* (Caryophyllaceae) and *Leymus mollis* (Poaceae) spatial segregation on subarctic coastal dunes.** *Am J Bot.* 2002; **89**(3): 479–485.

[PubMed Abstract](#) | [Publisher Full Text](#)

John UP, Polotnianka RM, Sivakumaran KA, *et al.*: **Ice recrystallization inhibition proteins (IRIPs) and freeze tolerance in the cryophilic Antarctic hair grass *Deschampsia antarctica* E. Desv.** *Plant Cell Environ.* 2009; **32**(4): 336–348.

[PubMed Abstract](#)

Krieger E, Joo K, Lee J, *et al.*: **Improving physical realism, stereochemistry, and side-chain accuracy in homology modeling: Four approaches that performed well in CASP8.** *Proteins.* 2009; **77**(Suppl 9): 114–122.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Middleton AJ, Marshall CB, Faucher F, *et al.*: **Antifreeze Protein from Freeze-Tolerant Grass Has a Beta-Roll Fold with an Irregularly Structured Ice-Binding Site.** *J Mol Biol.* 2012; **416**(5): 713–724.

[PubMed Abstract](#) | [Publisher Full Text](#)

Raymond J: **Ice structuring by ice-binding protein of *Leymus mollis***

DSCN5945.JPG. *figshare.* Figure. 2020a.

<http://www.doi.org/10.6084/m9.figshare.12401966.v1>

Raymond J: ***Leymus mollis* ice recrystallization inhibition.** *figshare.* Figure. 2020b.

<http://www.doi.org/10.6084/m9.figshare.12401954.v1>

Raymond J: **PCR amplification of IBP and 18S genes of *Leymus mollis*.** *figshare.* Figure. 2020c.

<http://www.doi.org/10.6084/m9.figshare.12401957.v1>

Raymond JA, Fritsen CH: **Semipurification and ice recrystallization inhibition activity of ice-active substances associated with Antarctic photosynthetic organisms.** *Cryobiology.* 2001; **43**(1): 63–70.

[PubMed Abstract](#) | [Publisher Full Text](#)

Sandve SR, Rudi H, Asp T, *et al.*: **Tracking the evolution of a cold stress associated gene family in cold tolerant grasses.** *BMC Evol Biol.* 2008; **8**: 245.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sidebottom C, Buckley S, Pudney P, *et al.*: **Heat-stable antifreeze protein from grass.** *Nature.* 2000; **406**(6793): 256.

[PubMed Abstract](#) | [Publisher Full Text](#)

Waterhouse A, Bertoni M, Bienert S, *et al.*: **SWISS-MODEL: homology modelling of protein structures and complexes.** *Nucleic Acids Res.* 2018; **46**(W1): W296–W303.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status: 

Version 1

Reviewer Report 20 July 2020

<https://doi.org/10.5256/f1000research.26842.r65703>

© 2020 Davies P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Peter L. Davies 

Protein Function Discovery Group, Department of Biomedical and Molecular Sciences, Queen's University, Kingston, ON, Canada

Sformo and Raymond have prepared aqueous extracts of a hardy coastal grass (*Leymus mollis*) collected in Alaska and have observed the presence of an ice-binding protein (IBP) based on ice crystal faceting and the inhibition of ice recrystallization. From another extract they prepared DNA and used primers based on conserved regions in other grass IBP transcripts to amplify the *L. mollis* gene, which fortunately lacks introns. From the sequence of this DNA they were able to provide the amino acid sequence of the IBP domain involved in ice recrystallization. They used this to model the three-dimensional structure of the domain, which looks very much like that of *Lolium perenne*. Another set of primers was used to amplify a diagnostic section of the 18S ribosomal RNA gene whose sequence identified the species.

There is an element of comparative science here, and one would not need to see a repeat of this detailed analysis on other grasses/cereals. The presence of sequence homologs in transcriptomes as a way of seeing the distribution and variation of this IBP type should suffice from here on. Nevertheless, it is interesting to see this grass from a high latitude in Alaska (Beaufort Sea coast) extracted and analyzed. This report is also a good illustration of the use of molecular modeling to avoid solving a structure that would not differ much from the template. Indeed, the main features of the *Lolium perenne* IBP fold are retained in the *L. mollis* model, including the outward bulge of the beta-solenoid at one end on the non-ice-binding side.

Suggestions for improvement:

- The Abstract should include the percent identity of the IRI domain between the two grass species. This type of information should also be provided in the Introduction where the sequences of the IBPs of *L. perenne* and *D. antarctica* are compared. It is not helpful to say their sequences “are very similar”. In the last paragraph of the Introduction the *L. mollis* IBP sequence is again said to be “very similar” to those of the other two grasses. Please provide % identity and % similarity.
- Figure 1A shows ice structuring. However, since panel B also shows ice crystals it is important to point out to the average reader that the distinction lies in the faceting seen in A where flat, angular

surfaces are the result of IBP binding to ice. The ice in B is smoother and rounded.

- Figure 2 is a rather poor presentation of ice recrystallization. One must look closely at the ice control images to see the grains are larger after 21 h. The methods say the ice recrystallization was done in water. From our experience the inclusion of 150 mM NaCl makes a huge difference to the visibility of the ice grains. The other suggestion is to take the starting picture at zero time with the ice held colder than -3°C . We see substantial recrystallization happening in the first hour, even at -6°C , and will typically hold the ice wafer at -10°C until the starting picture is captured, and then raise the temperature for more rapid recrystallization. I recommend that this analysis be repeated and improved.
- In Methods, under 'Structure prediction', it is stated that three structures were proposed by SWISS MODEL operating with *L. perenne* IBP as a template. Then, the one with the 'highest sequence identity' was selected. Surely the sequence identity is the same for all models because they are based on the same template. We would typically compare models by root mean square deviation (RMSD) – how the atoms deviate from the model on average. Is this what is meant here?
- Although the sequence of the *L. mollis* open reading frame was incomplete, it would be useful to report the sequence upstream of the ice-binding domain. Presumably this is a series of leucine-rich repeats similar to those seen in other grass IBPs?
- In reference to ice crystal axes the a- and c- should be italicized.
- Replace the contraction "didn't" with "did not".

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Protein biochemistry, structural biology, ice-binding proteins

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research