



SYSTEMATIC REVIEW

REVISED Application of genus *Cassia* in the treatment of Constipation: A systematic review [version 2; peer review: 1 approved, 1 approved with reservations]

Muhammad Shahzad Aslam^{1,2}

¹School of Traditional Chinese Medicine, Xiamen University Malaysia, Sepang, Selangor, 43900, Malaysia

²Department of Chemistry, Xiamen University Malaysia, Sepang, Selangor, 43900, Malaysia

V2 First published: 05 Mar 2019, 8:256
<https://doi.org/10.12688/f1000research.17893.1>
 Latest published: 08 Mar 2021, 8:256
<https://doi.org/10.12688/f1000research.17893.2>

Abstract

Purpose: Role of genus *Cassia* in the treatment of Constipation
Methods: Methodological analysis, systematic review, and meta-analysis of identified studies using RevMan
Result and Discussion: *Cassia fistula* was partially effected in treating constipation however there is a need for improvement in the protocol of studies to reduce biases. These results were only limited to one species so it cannot be generalized among all species of *Cassia*.
Conclusion: *Cassia fistula* is partially effective in reducing the pain and consistency of stool during constipation among children.

Keywords

Cassia, *Cassia fistula*, Constipation, Pediatric gastroenterology

Open Peer Review

Approval Status

	1	2
version 2 (revision) 08 Mar 2021	 view	
version 1 05 Mar 2019	 view	 view

1. **Massimo Bellini**, University of Pisa, Pisa, Italy
2. **Farhad Shokraneh** , University of Nottingham, Nottingham, UK

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Muhammad Shahzad Aslam (aslam.shahzad@xmu.edu.my)

Author roles: **Aslam MS:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This research was supported by Xiamen University Malaysia.
The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Aslam MS. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Aslam MS. **Application of genus *Cassia* in the treatment of Constipation: A systematic review [version 2; peer review: 1 approved, 1 approved with reservations]** F1000Research 2021, 8:256 <https://doi.org/10.12688/f1000research.17893.2>

First published: 05 Mar 2019, 8:256 <https://doi.org/10.12688/f1000research.17893.1>

REVISED Amendments from Version 1

Typing mistakes have been corrected.

Any further responses from the reviewers can be found at the end of the article

Introduction

Constipation is a clinical disorder attributed to ineffectual colonic impulsion and/or increased resistance to the proliferation of colonic matters¹. Approximately 20% of the world population suffers from chronic constipation². It is one of the most common pediatric problems³. It was found to be the second most stated disorder in the field of pediatric gastroenterology. Treatment costs for children with constipation will be around three times higher than children without constipation in the United States⁴. African American children, particularly girls, are greatly affected by constipation, which has been associated with poor hygiene conditions⁵.

Commonly, constipation is treated by Cisapride in children⁶, other treatments include polyethylene glycol 3350 and lactulose, however polyethylene glycol 3350 has been found to be more effective⁷. Supplemented and non-supplemented yogurt helps in reducing abdominal pain and to enhance defecation frequency⁸. It has been observed that different species of *Cassia* act effective as a laxative such as *Cassia fistula*, *Cassia alata*, and *Cassia augustifolia*^{9–11}. The genus *Cassia* is well known in alternative medicine as hepatoprotective¹², laxative, and in the treatment of ringworm infection¹³, skin diseases¹⁴ and leprosy¹⁵. It has many pharmacological properties including acting as a hypolipidemic agent¹⁶, anti-microbial¹⁷, anti-fungal¹⁸, and anti-cancer agent¹⁹. Genus *Cassia* contains a number of bioactive compounds such as anthraquinone²⁰, tannin²¹, coumarins²², triterpene, volatile oil²³, phenolic glycoside²⁴, flavonoids²⁵ from different parts of the plant. Different species of *Cassia* possess laxative properties due to various anthraquinone derivative such as aloe-emodin, rhein, chrysophanol and chrysoobtusin. In this review, we systematically assessed the laxative potential of different species of *Cassia*.

Methods

Literature search strategy

A systematic literature search was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. Using the keywords (Senna AND Laxatives AND Clinical trial) (Cassia AND Laxatives AND Clinical trial) (Cassia AND Clinical trial) (Senna AND Clinical trial) and publication range from 01 January 1960 till 31 December 2018 for identification of the records. **Table 1** shows the search strategy for PubMed Central. During screening of the records only full-length open access articles were considered. Abstract only or closed access articles were excluded¹¹. Only articles involving children aged between 2–15 were included. All review articles, *in-vivo* studies and those >10 years from the search date were excluded. A preliminary search of the PubMed, CNKI, Scopus, Web of Science, Google Scholar and PsycINFO databases and digital archive such as PubMed Central, yielded 2207 papers published in English from the last ten years. Duplicate and irrelevant articles were removed (n=2203). One article was further removed during screening due to closed access (n=3). One publication was removed because the article did not meet the eligibility criteria (n=2). A PRISMA Flow Diagram is given in **Figure 1**.

Literature screening

Identification of articles were performed at level 1 using the search strategy as mentioned in **Table 1**. Duplicate articles, irrelevant articles such as polyherbal formulation, review articles, or any article other than *Cassia* or *Senna* were removed at level 2. Only four⁴ articles were identified as being relevant. One record was excluded due to not being a full text article. Abstracts were being reviewed for the following inclusion and exclusion criteria at level 4 and one article was removed for not meeting the eligibility criteria i.e. Randomized, clinical trial on Constipation, full-length open access articles, Pediatric Functional Constipation (age range: 2–15 years).

Eligibility criteria

Types of studies. The author has selected studies of randomized open label, prospective, controlled, parallel-group clinical trial for meta-analysis. Baseline characteristics of randomized trials

Table 1. Search strategy.

PubMed Central	
("cassia"[MeSH Terms] OR "cassia"[All Fields]) AND ("clinical trial"[All Fields] OR "clinical trials as topic"[MeSH Terms] OR "clinical trial"[All Fields]) AND ("1960/01/01"[PubDate]: "2018/12/31"[PubDate])	((("cassia"[MeSH Terms] OR "cassia"[All Fields]) AND ("laxatives"[All Fields] OR "laxatives"[MeSH Terms] OR "laxatives"[All Fields])) AND ("clinical trial"[All Fields] OR "clinical trials as topic"[MeSH Terms] OR "clinical trial"[All Fields]) AND ("1960/01/01"[PubDate]: "2018/12/31"[PubDate])
("Senna"[MeSH Terms] OR "Senna"[All Fields]) AND ("clinical trial"[All Fields] OR "clinical trials as topic"[MeSH Terms] OR "clinical trial"[All Fields]) AND ("1960/01/01"[PubDate]: "2018/12/31"[PubDate])	((("senna plant"[MeSH Terms] OR ("senna"[All Fields] AND "plant"[All Fields]) OR "senna plant"[All Fields] OR "senna"[All Fields]) AND ("laxatives"[All Fields] OR "laxatives"[MeSH Terms] OR "laxatives"[All Fields])) AND ("clinical trial"[All Fields] OR "clinical trials as topic"[MeSH Terms] OR "clinical trial"[All Fields]) AND ("1960/01/01"[PubDate]: "2018/12/31"[PubDate])




Figure 1. PRISMA flow diagram.

of studies included on pediatric functional constipation are presented in [Table 4](#). Characteristics of the studies included are mentioned in [Table 5](#).

Types of participants. The author included studies involving patients (aged 2–15 years) with Functional constipation. The diagnosis of Functional constipation was according to the Rome III criteria²⁶. Inclusion and exclusion criteria were based on Study design, participants, intervention, outcome (SPIO) criteria and indicated in [Table 2](#).

Types of interventions. Included studies were focused on the role Cassia in the treatment of Functional constipation. Unfortunately, there were only two studies identified.

Types of outcomes. Eligible studies included consisted of the following outcomes: improvement in the episodes of


fecal incontinence per week, improvement in the episodes of retentive posturing per week, improvement in the average of severity of pain of defecation (by VAS), improvement in defecation frequency per week, patient's drug compliance and improvement in the average of consistency of stool defecated (by VAS).

Methodological quality assessment (MQ)

Methodological quality assessment was made on the basis of following criteria. 1) Aims and Hypothesis clearly defined, adequate sample representation, patient care quality, ethical approval protocol, outcomes assessment, validity and reliability of outcome measure, attempt to blind researcher, follow-up, appropriate statistical analysis and missing data reported. Ten item defined evaluation of methodological quality (MQ) is presented in [Figure 2](#). Risk of Bias were assessed using Cochrane collaboration's tool on the basis of the following criteria such as

Table 2. Study design; participants; intervention; outcome (SPIO) criteria.

	Inclusion Criteria	Exclusion Criteria
Study design	Randomized, clinical trial on Constipation, full-length open access articles, Pediatric Functional Constipation	All review articles, irrelevant articles, Exclude abstract only articles, clinical trial on adults.
Participants	Children (age range: 2–15 years)	Adults
Intervention	<i>Cassia fistula</i> was delivered to Pediatric with Functional Constipation	
Outcomes	Role of the <i>Cassia fistula</i> emulsion in Pediatric Functional Constipation	

**Figure 2. Methodological quality assessment of the 2 studies included in the meta-analysis (0=No/not reported, 1=Yes).**

selection bias, performance bias, attrition bias, reporting bias and miscellaneous. Cochrane Collaboration's tool for assessing the risk of bias was used and the results are presented in Table 6²⁷.

Data extraction

The following data were extracted according to study characteristics (e.g., first author, year of publication, search dates, and

number of included studies), patient characteristics (functional constipated children, aged between 2–15), sample size, study type (e.g., Randomized open label, prospective, controlled, parallel-group clinical trial study), randomization methods (e.g., “systematic randomization and simple randomization”) and outcome measures/variables (e.g., improvement in defecation frequency per week, improvement in the episodes of fecal incontinence per week, improvement in the episodes of

retentive posturing per week, improvement in the average of severity of pain of defecation (by VAS), improvement in the average of consistency of stool defecated (by VAS) and patient's drug compliance). Data extraction was performed by Muhammad Shazad Aslam. Transcripts were analysed, coded and data was extracted using the demo version of qualitative data analysis software *Atlas.ti 8.0*. **Table 3** represent all the data that was extracted. All the meta-data are available as Dataset 1.

Statistical analysis

Meta-analysis was conducted using the *Review Manager (RevMan) 5.3* software²⁸. The summary measures were reported as odds ratios (ORs) or as a standard mean difference (SMD) with 95% confidence intervals (CI). The presence of heterogeneity among trials was assessed using the Chi-square test, and the extent of the inconsistency was measured by I² statistics. Output file from RevMan is available as Underlying data²⁹.

Results and discussion

Both of the selected studies were not blinded during intervention and outcome assessment that will result in performance bias

and detection bias respectively. These biases occur where the investigators know about the participant's treatment group. Performance bias can also refer to the fact that participants can change their responses or behaviour if they know which group they are allocated in. Blinding of outcome assessment may decrease the risk of the investigator or participant being aware of the treatment that a patient is receiving. If the participants and the caregivers are aware of the intervention and outcome that may affect the behavior of the participants, these behavioral changes may affect the performance of the treatment. Clinical trials on adults were also excluded, such as a randomized clinical trial of a phytotherapeutic compound containing *Pimpinella anisum*, *Foeniculum vulgare*, *Sambucus nigra*, and *Cassia augustifolia* for chronic constipation⁹. Results of both included studies were non-significant when comparing their baseline characteristics of pediatric functional constipation as presented in **Table 5**. During analysis of study characteristics, it was found that both of studies demonstrated *Cassia fistula* is helping to treat constipation among the children as shown in **Table 3**, but there is a risk of bias according to Cochrane Collaboration's tool (**Table 6**). Moreover, both studies found were from one country (Iran). During a methodological assessment, many flaws were identified

Table 3. Data extraction.

Characteristics	Seyyed Ali Mozaffarpur	Mohammad Reza Esmaeilidooki
Publication Year	2012	2016
Type of Study	Randomized, clinical trial	This randomized open label, prospective, controlled, parallel-group clinical trial study
Age	Age between 4–13 years	Aged between 2 – 15 years
Randomization	systematic randomization	Simple randomization
Total Sample Size	81	51
Variables	Frequency of defecation, consistency of stools, and severity of pain during defecation, retentive posturing and fecal incontinence per week	Improvement in defecation frequency per week, improvement in the episodes of fecal incontinence per week, improvement in the episodes of retentive posturing per week, improvement in the average of severity of pain of defecation (by VAS), improvement in the average of consistency of stool defecated (by VAS) and patient's drug compliance
Length of each contact with the participant/ caregivers	Clinical efficacy and tolerance were assessed using weekly sheets, parents completed every night. They were given three sheets (included seven questions in seven columns) to complete them daily for 3 weeks.	During the study, we had regular phone calls with the parents to check the probable complications, treatment (taking the prescribed drugs) and data filling process. If there were any serious questions or problems, we visited the child. At the end of 4 weeks of treatment, the children were visited and the filled out forms were taken and evaluated.
Blinding of Experiment	The investigators, the children and their parents were aware of the study group assignment.	Fortunately, due to the developing socioeconomic conditions of the people in these regions in recent years, we were able to keep in touch with all the patients during the study period by phone call.
Diagnosis	Rome III criteria of functional constipation	Rome III criteria of functional constipation
Diagnostic test	Paraclinics like anorectal manometry, thyroid function tests, anti-tTG, and other laboratory tests	thyroid function tests, anti-tTG, and etc. If it remained any doubt, barium study and anorectal manometry would be performed

Table 4. Baseline characteristics of randomized trials of studies included in pediatric functional constipation.

Variable in Treatment (T)	Mozaffarpur, 2012 (T)	Esmailidooki, 2016 (T)	Mozaffarpur, 2012 (C)	Esmailidooki, 2016 (C)	Variable in Control (C)
Sample size in Treatment (N)	41	52	40	57	Sample size in Control (N)
Age, months, Mean(\pm SD)	69.4(\pm 24.3)	64.6(\pm 25.2)	65.9(\pm 19.1)	55.2(\pm 31.2)	Age, months, Mean(\pm SD)
Sex, Male, (n) (%)	29(70.7%)	33	23(57.5%)	30	Sex, Male, (n) (%)
Weight, Kg Mean(\pm SD)	21.7 (\pm 7.2)	20.5(\pm 7.2)	20.7(\pm 7.8)	18.5(\pm 8.9)	Weight, Kg(\pm SD)
Duration of constipation, months, Mean(\pm SD)	34.2(\pm 25.9)	31.1(\pm 24.6)	30.8(\pm 22.8)	23.5(\pm 21.8)	Duration of constipation, months, Mean(\pm SD)
Defecation \leq 2 per week, n (%)	32(78%)	41	30(75%)	52	Defecation \leq 2 per week, n (%)
Incontinence, n (%)	31(75.6%)	34	27(67.5%)	37	Incontinence, n (%)
History of previous treatment, n (%)	32 (78%)	43	28(70%)	51	History of previous treatment, n (%)
Retentive posturing, n (%)	32(78%)	40	29(72.5%)	37	Retentive posturing, n (%)

Table 5. Study characteristics.

Author, year	Study Design	Hypothesis	Statistical analysis	Software	Outcomes
Mozaffarpur, 2012	Randomized, clinical trial	The author hypothesized that <i>Cassia fistula</i> emulsion (CFE) would be as effective or better than Mineral oil (MO) in treating FC.	The statistical analyses included the determination of means and Standard deviation (SDs), t test, χ^2 test, ANOVA repeated measures and Fisher's exact test, with significance accepted at the 5% level.	SPSS (version 17),	CFE was most effective than MO in the 3-week treatment of children with FC.
Esmailidooki, 2016	Randomized, clinical trial	N/A	The statistical analyses included the determination of means and SDs, t test, χ^2 test, ANOVA repeated measures and Fisher's exact test, with significance accepted at the 5% level.	SPSS IBM20 and STATA 11.2	Significant improvement when compared with the control group however unable to find substantial evidence of the role of identified bioactive compounds due to limitation as it requires further investigation

Table 6. Cochrane Collaboration's tool for assessing risk of bias.

Study (author, year)	Selection bias	Performance bias	Detection bias	Attrition bias	Reporting bias	Other bias
Mozaffarpur, 2012	Unclear	Yes	Yes	No	No	Unclear
Esmailidooki, 2016	Unclear	Yes	Yes	No	No	Unclear

such as inadequate patient care, attempt to blind the researcher and missing data (Figure 2). During meta-analysis, the comparison was made before and after treatment among different variables such as defaecation, fecal incontinence, retentive posturing, the severity of pain, and consistency of stool. All the variable (before and after treatment) were found to be symmetrical when plotted on a funnel plot as shown in Figure 4, Figure 6, Figure 8, Figure 10, Figure 12, and Figure 14 respectively. The overall effect for some variables is statistically insignificant (P=0.11, P=0.49, P=0.24) such as fecal incontinence, retentive posturing, and acceptance, tolerance respectively. High heterogeneity was found in two variables i.e severity of pain (90%) and consistency of stool (77%). All the forest plot of defaecation, fecal incontinence, retentive posturing, severity of pain,

consistency of stool and acceptance and tolerance are represented in Figure 3, Figure 5, Figure 7, Figure 9, Figure 11 and Figure 13 respectively.

Conclusion

After evaluation of results, it was found that *Cassia fistula* was not completely effective. It was partly effective in reducing the pain and consistency of stool during constipation. However, these results cannot be generalized among all population. A well designed, expert validated protocol is required in the future. There is a need to develop an instrument that will be free from bias. Moreover, the results cannot be generalized among all species of *Cassia* as the studies available are only for one species. There is a need to isolate identified bioactive compounds from




Figure 3. Forest plot in defaecation before and after treatment.




Figure 4. Funnel plot showing overall standardized mean difference in defaecation before and after treatment.





Figure 5. Forest plot in fecal incontinence before and after treatment.




Figure 6. Funnel plot showing overall standardized mean difference in fecal incontinence before and after treatment.





Figure 7. Forest plot in retentive posturing before and after treatment.




Figure 8. Funnel plot showing overall standardized mean difference in retentive posturing before and after treatment.





Figure 9. Forest plot in severity of pain before and after treatment.




Figure 10. Funnel plot showing overall standardized mean difference in severity of pain before and after treatment.




Figure 11. Forest plot in consistency of stool before and after treatment.




Figure 12. Funnel plot showing overall standardized mean difference in consistency of stool before and after treatment.




Figure 13. Forest plot in acceptance and tolerance after treatment.




Figure 14. Funnel plot showing overall standardized mean difference in acceptance and tolerance after treatment.

different species of *Cassia* and evaluate the effect of different factors such as duration of constipation, defecation, incontinence or retentive posturing under clinical trial.

Declarations

Data availability

Underlying data. Open Science Framework: Application of genus *Cassia* in the treatment of Constipation: A systematic review. <https://doi.org/10.17605/OSF.IO/PKR4N>²⁹

This project contains the following underlying data:

- *Cassia Senna* for Constipation.rm5 (study RevMan file)

- Quotation Manager.xlsx (study characteristics of citations included in this study)

Reporting guidelines

Open Science Framework: PRISMA diagram and flowchart for the study “Application of genus *Cassia* in the treatment of constipation: A systematic review”. <https://doi.org/10.17605/OSF.IO/PKR4N>²⁹

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

References

- Bharucha AE: **Constipation.** *Best Pract Res Clin Gastroenterol.* 2007; **21**(4): 709–731.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Camilleri M, Ford AC, Mawe GM, *et al.*: **Chronic constipation.** *Nat Rev Dis Primers.* 2017; **3**: 17095.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Le Luyer B, Ménard M: **[Constipation in children].** *Rev Prat.* 1998; **48**(4): 376–381.
[PubMed Abstract](#)
- Liem O, Harman J, Benninga M, *et al.*: **Health utilization and cost impact of childhood constipation in the United States.** *J Pediatr.* 2009; **154**(2): 258–262.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Uc A, Hyman PE, Walker LS: **Functional gastrointestinal disorders in African American children in primary care.** *J Pediatr Gastroenterol Nutr.* 2006; **42**(3): 270–274.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nurko S, Garcia-Aranda JA, Worona LB, *et al.*: **Cisapride for the treatment of constipation in children: A double-blind study.** *J Pediatr.* 2000; **136**(1): 35–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Gremse DA, Hixon J, Crutchfield A: **Comparison of polyethylene glycol 3350 and lactulose for treatment of chronic constipation in children.** *Clin Pediatr (Phila).* 2002; **41**(4): 225–229.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Guerra PV, Lima LN, Souza TC, *et al.*: **Pediatric functional constipation treatment with *Bifidobacterium*-containing yogurt: a crossover, double-blind, controlled trial.** *World J Gastroenterol.* 2011; **17**(34): 3916–3921.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Picon PD, Picon RV, Costa AF, *et al.*: **Randomized clinical trial of a phytotherapeutic compound containing *Pimpinella anisum*, *Foeniculum vulgare*, *Sambucus nigra*, and *Cassia augustifolia* for chronic constipation.** *BMC Complement Altern Med.* 2010; **10**: 17.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mozaffarpur SA, Naseri M, Esmailidooki MR, *et al.*: **The effect of *cassia fistula* emulsion on pediatric functional constipation in comparison with mineral oil: a randomized, clinical trial.** *Daru.* 2012; **20**(1): 83.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Thamlikitkul V, Bunyapraphatsara N, Dechatiwongse T, *et al.*: **Randomized controlled trial of *Cassia alata* Linn. for constipation.** *J Med Assoc Thai.* 1990; **73**(4): 217–222.
[PubMed Abstract](#)
- Jafri MA, Jalis Subhani M, Javed K, *et al.*: **Hepatoprotective activity of leaves of**

- Cassia occidentalis* against paracetamol and ethyl alcohol intoxication in rats.** *J Ethnopharmacol.* 1999; **66**(3): 355–361.
[PubMed Abstract](#) | [Publisher Full Text](#)
13. Abubacker MN, Ramanathan R, Senthil Kumar T: **In vitro antifungal activity of *Cassia alata* linn. Flower extract.** *Indian Journal of Natural Products and Resources.* 2008; **7**(1): 6–9.
[Reference Source](#)
 14. Benjamin TV, Lamikanra A: **Investigation of *Cassia alata*, a Plant Used in Nigeria in the Treatment of Skin Diseases.** *Q J Crude Drug Res.* 1981; **19**(2–3): 93–96.
[Publisher Full Text](#)
 15. Singh S, Singh SK, Yadav A: **A Review on Cassia species: Pharmacological, Traditional and Medicinal Aspects in Various Countries.** *American Journal of Phytomedicine and Clinical Therapeutics.* 2013; [cited 2019 Jan 11]; **3**: 291–312.
[Reference Source](#)
 16. Patil UK, Saraf S, Dixit VK: **Hypolipidemic activity of seeds of *Cassia tora* Linn.** *J Ethnopharmacol.* 2004; **90**(2–3): 249–252.
[PubMed Abstract](#) | [Publisher Full Text](#)
 17. Gaddam SA, Kotakadi VS, Sai Gopal DV, et al.: **Efficient and robust biofabrication of silver nanoparticles by *Cassia alata* leaf extract and their antimicrobial activity.** *J Nanostruct Chem.* 2014; **4**(1): 82.
[Publisher Full Text](#)
 18. Sumathy V, Zakaria Z, Jothy SL, et al.: **In vitro and in vivo antifungal activity of *Cassia surattensis* flower against *Aspergillus niger*.** *Microb Pathog.* 2014; **77**: 7–12.
[PubMed Abstract](#) | [Publisher Full Text](#)
 19. Yuenyongsawad S, Bunluepuech K, Wattanapiromsakul C, et al.: **Anti-cancer activity of compounds from *Cassia garrettiana* heartwood.** *Songklanakarinn J Sci Technol.* 2014; **36**(2): 189–194, [cited 2019 Jan 11]
[Reference Source](#)
 20. Jung HA, Ali MY, Jung HJ, et al.: **Inhibitory activities of major anthraquinones and other constituents from *Cassia obtusifolia* against β -secretase and cholinesterases.** *J Ethnopharmacol.* 2016; **191**: 152–160.
[PubMed Abstract](#) | [Publisher Full Text](#)
 21. Sivakumar V, Ilanhtiraiyan S, Ilayaraja K, et al.: **Influence of ultrasound on Avaram bark (*Cassia auriculata*) tannin extraction and tanning.** *Chem Eng Res Des.* 2014; **92**(10): 1827–1833.
[Publisher Full Text](#)
 22. Ngoc TM, Nhiem NX, Khoi NM, et al.: **A new coumarin and cytotoxic activities of constituents from *Cinnamomum cassia*.** *Nat Prod Commun.* 2014; **9**(4): 487–488.
[PubMed Abstract](#)
 23. Jeyaratnam N, Nour AH, Kanthasamy R, et al.: **Essential oil from *Cinnamomum cassia* bark through hydrodistillation and advanced microwave assisted hydrodistillation.** *Ind Crops Prod.* 2016; **92**: 57–66.
[Publisher Full Text](#)
 24. Zeng J, Xue Y, Lai Y, et al.: **A new phenolic glycoside from the barks of *Cinnamomum cassia*.** *Molecules.* 2014; **19**(11): 17727–17734.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 25. Luximon-Ramma A, Bahorun T, Soobrattee MA, et al.: **Antioxidant activities of phenolic, proanthocyanidin, and flavonoid components in extracts of *Cassia fistula*.** *J Agric Food Chem.* 2002; **50**(18): 5042–5047.
[PubMed Abstract](#) | [Publisher Full Text](#)
 26. Fabricant DS, Farnsworth NR: **The value of plants used in traditional medicine for drug discovery.** *Environ Health Perspect.* 2001; **109** Suppl 1: 69–75.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 27. Collaboration C: **Cochrane handbook for systematic reviews of interventions.** 2008.
[Reference Source](#)
 28. Collaboration C: **Review manager (RevMan)[computer program].** 2011.
 29. Aslam MS: **Application of genus *Cassia* in the treatment of Constipation: A systematic review.** 2019.
<http://www.doi.org/10.17605/OSF.IO/PKR4N>

Open Peer Review

Current Peer Review Status:  

Version 2

Reviewer Report 06 May 2021

<https://doi.org/10.5256/f1000research.55132.r82025>

© 2021 Bellini M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Massimo Bellini

Gastrointestinal Unit, Department of Translational Research and New Technologies in Medicine and Surgery, University of Pisa, Pisa, Italy

I've no further particular methodological concerns. My main concern is still the same: I'm really puzzled about the usefulness of a review which takes into consideration only two papers and altogether only 132 patients.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Gastroenterology: Functional Digestive Disorders.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 28 June 2019

<https://doi.org/10.5256/f1000research.19569.r50310>

© 2019 Shokrane F. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Farhad Shokrane

University of Nottingham, Nottingham, UK

The author investigates the effect of *Cassia fistula* on three outcomes based on two included

studies. It seems a valuable research that could potentially direct the pharmaceutical companies toward a new agent to treat the constipation.

Title

The title should be "Cassia fistula for treatment of children and adolescents with constipation: A systematic review of randomized controlled trials".

Authorship

The systematic review requires a process that involves at least two people for screening and data extraction. I suggest the author to find another systematic reviewer to be involved in this work.

Abstract

The abstract is very brief and sentences are incomplete. Abstract should be re-written following the example from similar systematic reviews and following [PRISMA for Abstract](#).

Minor edits

1. "act an effective as a laxative" should be "act as effective as laxatives".
2. "search data" should be "search date".

Methods

1. The standard search filter for finding RCTs as reported in Cochrane Handbook should be used.
2. It is advised in Cochrane Handbook searching Embase, MEDLINE/PubMed, and CENTRAL for systematic reviews of RCTs and this review reports searching only MEDLINE/PubMed.
3. Excluding closed access papers without trying to order them from library or contacting the authors introduces full-text or open access bias.
4. The structure of the method should change and start with 1: Eligibility Criteria 2: Databases and search strategy 3: Screening process 4: Data extraction process 5: Assessment of risk of bias process 6: Data analysis plan. Currently, the first section of the methods reports info on search and eligibility criteria while there are specific subheadings for these sections.
5. The rationale behind excluding studies older than 10 years from search date is not clear.
6. The reported number of search results 2207 is not reasonable. A proper search should not find more than 100-200 results for this topic.
7. The search date has not been reported.
8. Following a published Cochrane review of RCTs in terms of structure and all the items in [PRISMA Checklist for full systematic reviews](#) is recommended.
9. Constipation is missing from search strategy and MeSH terms in the table.
10. The numbers in PRISMA flow diagram does not match the numbers reported in the text and it does not seem to be correct. Following one of the existing published systematic reviews of

RCTs from Systematic Reviews journal or Cochrane Database of Systematic Reviews is recommended.

11. The quality of included studies is being assessed using Cochrane Risk of Bias tool. The role of MQ and using Atlas.ti is not clear in this review. These two are not usually used for SR of RCTs.
12. Studying the meta-analysis section in the Cochrane Handbook which is freely available online could help revising the statistical analysis.
13. Table 2 should report PICOS not SPIO.
14. The time point for the outcome is not clear.
15. The outcome measure has not been described properly.
16. The complications and adverse effects have not been reported in the review.

Results

1. The sample size reported in the Data Extraction table does not match the numbers in meta-analysis. For example the total for experimental and control group in Review Manager file is 109 for Esmailidooki 2016 and 71 for Mozaffarpur 2012 while in the table is 51 and 81 respectively.
2. The analysis for before treatment is not required. All before treatments can be deleted.
3. Funnel plots only work with about 10-20 studies. All funnel plots can be deleted.
4. Risk of bias based on Cochrane tool is not clear. It should be low risk, unclear risk and high risk. The current description of Yes/No does not provide enough information.
5. In meta-analysis, "one randomized always analysed" while the numbers in Table 4 are not the numbers in meta-analysis in Review Manager.
6. The control groups in two studies are different so I am not sure if combining the data from two studies in a meta-analysis is a right when there is such heterogeneity in PICOS components.
7. SD is greater than mean is some outcomes which is an indicator of skewed data or error in reporting. Contacting the trials authors might make some of the points clear.

Discussion and Conclusion

Current data and status of the review may not be appropriate for discussion or making a conclusion. The current evidence is limited and more rigorous studies will be required to make any conclusion or recommendations.

I understand the author's passion for conducting a systematic review and so I request the author to work with an expert supervisor in the field before revising the work.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Partly

Are sufficient details of the methods and analysis provided to allow replication by others?

Partly

Is the statistical analysis and its interpretation appropriate?

No

Are the conclusions drawn adequately supported by the results presented in the review?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Evidence synthesis, Systematic review, Overview, Scoping review, Rapid review, Randomized controlled trials, Open data, Open access, Open source, Medical journalism, Peer-review, Open source, Outcomes, Interventions, Cochrane

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 02 Mar 2021

Muhammad Shahzad Aslam, Xiamen University Malaysia, Malaysia

WHO defines '**Adolescents**' as individuals in the 10-19 years. So title cannot be changed

Competing Interests: No

Reviewer Report 23 May 2019

<https://doi.org/10.5256/f1000research.19569.r47237>

© 2019 Bellini M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Massimo Bellini

Gastrointestinal Unit, Department of Translational Research and New Technologies in Medicine and Surgery, University of Pisa, Pisa, Italy

I'm really puzzled about the usefulness of a review which takes into consideration only two papers and altogether only 132 patients. Constipation, as the author himself states, is a very frequent condition, so the conclusions are inevitably flawed by this bias. The author should address this issue in the conclusions

Hence, **I have significant reservations.**

Moreover, [here](#) enclosed you'll find a copy of the paper. The authors can find some suggestions, as tracked changes and comments to the original manuscript, which could improve the quality of the manuscript.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Gastroenterology: Functional Digestive Disorders.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Response 23 May 2019

Muhammad Shahzad Aslam, Xiamen University Malaysia, sepang, Malaysia

Dear,

I have read the report. The selected studies were according to the criteria given inside the paper. So, there were only two selected studies for the systematic review. The comments you have mentioned are appreciable. Thanks

Competing Interests: No

Author Response 02 Mar 2021

Muhammad Shahzad Aslam, Xiamen University Malaysia, Malaysia

Please see inclusion criteria (Only articles involving children aged between 2–15 were included. Only articles involving children aged between 2–15 were included.)

Competing Interests: No

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research