


## Supplementary material

### The Human Release Hypothesis for biological invasions: human activity as a determinant of the abundance of invasive plant species

Heike Zimmermann\*, Patric Brandt, Joern Fischer, Erik Welk, Henrik von Wehrden

\* Corresponding author: heike.zimmermann@uni-leuphana.de



**Figure S1. *Rosa rubiginosa* (sweetbriar rose) occurrences used in our MAXENT model.**

Occurrences were taken from existing literature (see Appendix 2 in Supplementary material) and our field observations; blue = native occurrences; pink = invasive occurrences. For the final MAXENT model we included a randomly reduced dataset until data points were evenly distributed and no spatial autocorrelation was detected in the model residuals.

## Appendix 1

### Methods used for climatic envelopes and for niche equivalency test

Sweetbriar rose (*Rosa rubiginosa* L.) occurrences were taken from the literature (see Appendix 2 in Supplementary material) as well as from personal observations. In total, we collated 1425 occurrence records from the invasive range (Australia, New Zealand, North America, South America, South Africa) and 12132 occurrences from the native range (Europe, Asia, North Africa; see Figure S1 in Supplementary material and dataset 1). However, we did not include all native occurrences into our final MAXENT model because of a strong spatial autocorrelation in the model residuals. Therefore, we randomly reduced the native occurrences until data points were evenly distributed based on the Moran's I coefficient; this point was reached when the dataset was reduced to 3033 native occurrences.

As a priori assumption we defined the global range of our study species which would later serve as background for pseudo-absences. Nine bioclimatic variables were chosen from the WorldClim Global Climate Data-set<sup>1</sup> to define the global range of our study species: annual mean temperature (bioclim 1), isothermality (bioclim 3), temperature seasonality (bioclim 4), mean temperature of warmest quarter (bioclim 10), mean temperature of coldest quarter (bioclim 11), annual precipitation (bioclim 12), precipitation seasonality (bioclim 15), precipitation of the wettest quarter (bioclim 16), and precipitation of the driest quarter (bioclim 17). We followed the approach from VanderWal et al. (2009)<sup>2</sup> choosing the spatial extent that would provide the most accurate results and biologically meaningful fit. Therefore, we chose those bioclimatic variables that sufficiently outlined the potential global range of this species based on the total occurrences with the ArcGis<sup>®</sup> software and excluded redundant variables and monthly values. According to these criteria bioclimatic variables that highlighted almost the entire world as a potential habitat were neglected. All nine bioclimatic variables were added into one composite raster file, which was converted into a polygon representing the area with *R. rubiginosa* occurrences. Out of this area all bioclimatic variables were extracted to define the climatic envelope using the MAXENT (Maximum Entropy modelling for species geographic distributions) software (Vers. 3.3.3k)<sup>3,4</sup>. We chose five bioclimatic variables that contributed useful information by themselves: annual mean temperature (bioclim 1), mean diurnal range (bioclim 2), temperature annual range (bioclim 7), mean temperature of warmest quarter (bioclim 10), annual precipitation (bioclim 12), and precipitation seasonality (bioclim 15); and checked for collinearity with a correlation matrix.

Within the MAXENT software we selected a 20 % test percentage out of our dataset, 100 replicates (Bootstrap), and fade by clamping to mitigate clamping issues. Our climatic model based on the invasive occurrences had a high discrimination performance with an average test AUC of 0.90. Our climatic envelope model based on the native occurrences performed slightly less well with an average test AUC of 0.87. In both models, we detected no spatial autocorrelation in the residuals.


A threshold rule was applied to convert continuous suitability surfaces into binary outputs only showing areas that are suitable for modelled groups. We selected the threshold ‘maximum training sensitivity plus specificity logistic threshold’ which focuses on the correct classification of presences and background points<sup>5</sup>.

Moreover, we quantified the degree of niche overlap between the invasive and native niche by using the Schoener’s D index<sup>6,7</sup> from the R (v. 2.15) package ‘phyloclim’ (v. 0.9.2) which was recently shown to outperform other metrics when assessing niche overlaps<sup>8</sup>. Schoener’s D indicates the degree of accordance between two niches based on climate envelope model outputs and ranges from 0 = no overlap to 1 = full overlap. In a one-tailed test, Schoener’s D is compared to the percentiles of null distributions obtained from pseudo-replicated climate envelope models where localities from invasive and native populations are pooled and randomly partitioned. We set 100 as the factor for pseudo-replication which has been shown to be sufficient to reject the null hypothesis with high confidence<sup>7,9</sup>. The derived Schoener’s D of 0.31 ( $p < 0.0001$ ) verified the pattern observed in the map, that is, invasive and native populations occupied different climatic niches.


## References Appendix 1

1. Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. Very high resolution interpolated climate surfaces for global land areas. *Int J Climatol.* 2005; 25(15): 1965–78. Available from: <http://biogeo.berkeley.edu/>
2. VanDerWal J, Shoo LP, Graham C, Williams SE. Selecting pseudo-absence data for presence-only distribution modeling: How far should you stray from what you know? *Ecol Modell.* 2009; 220(4): 589–94. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0304380008005486>

3. Phillips SJ, Anderson RP, Schapire RE. Maximum entropy modeling of species geographic distributions. *Ecol Modell.* 2006; 190(3-4): 231–59. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S030438000500267X>
4. Elith J, Kearney M, Phillips S. The art of modelling range-shifting species. *Methods Ecol Evol.* 2010; 1(4): 330–42. Available from: <http://doi.wiley.com/10.1111/j.2041-210X.2010.00036.x>
5. Jiménez-Valverde A, Lobo J. Threshold criteria for conversion of probability of species presence to either–or presence–absence. *Acta Oecologica.* 2007; 31: 361–9. Available from: <http://www.sciencedirect.com/science/article/pii/S1146609X07000288>
6. Schoener T. The Anolis lizards of Bimini: resource partitioning in a complex fauna. *Ecology.* 1968; 49: 704–26. Available from: <http://www.jstor.org/stable/10.2307/1935534>
7. Warren DL, Glor RE, Turelli M. Environmental niche equivalency versus conservatism: quantitative approaches to niche evolution. *Evolution.* 2008; 62: 2868–83.
8. Rödder D, Engler JO. Quantitative metrics of overlaps in Grinnellian niches: advances and possible drawbacks. *Glob Ecol Biogeogr.* 2011; 20(6): 915–27. Available from: <http://doi.wiley.com/10.1111/j.1466-8238.2011.00659.x>
9. Broennimann O, Fitzpatrick MC, Pearman PB, Petitpierre B, Pellissier L, Yoccoz NG, et al. Measuring ecological niche overlap from occurrence and spatial environmental data. *Glob Ecol Biogeogr.* 2012; 21(4): 481–97.



**Figure S2. Histograms of the annual mean temperature [°C], mean diurnal range, temperature annual range and annual precipitation [mm] for all *Rosa rubiginosa* sites.** The annual mean temperature, mean diurnal range, temperature annual range and annual precipitation were the most important predictors contributing each more than 10% to the MAXENT models. The histograms show the density (frequency standardized to 100) of native (blue) and invasive (pink) occurrences for each parameter value (overlap in purple).



**Figure S3. Annual mean temperature [°C] and annual precipitation [mm] for selected invasive (pink) and native (blue) populations of *Rosa rubiginosa* (sweetbriar rose).** In the invasive range populations are located in Central (a, N=6) and Southern Argentina (b, N=7). In the native range populations are located in Spain (c, N=7) and Germany (d, N=11). Annual mean temperature and annual precipitation vary greatly within the invasive range and also within the native range. Smaller population sizes in the native range (mostly less than 10 individuals compared to hundreds of individuals in the invasive range) therefore are not explained by these two climatic variables.

## Appendix 2

**List of plant distribution data sources** for *Rosa rubiginosa* L. used by E. W. to compile the geographic distribution in the native and introduced areas. The sources are arranged by continents. All web based data sources were accessed in December 2012.

### North America

1. AKEPIC. *Alaska Exotic Plant Information Clearinghouse Database*. Available at: <http://aknhp.uaa.alaska.edu/maps/akepic/>
2. CalWeedMapper. *California Invasive Plant Council*. Available at: <http://calweedmapper.calflora.org/>
3. EDDMapS. *Early Detection and Distribution Mapping System*. Available at: <http://www.eddmaps.org/>
4. E-Flora BC. *Electronic Atlas of the Plants of British Columbia [eflora.bc.ca]*. Available at: <http://www.geog.ubc.ca/biodiversity/eflora/index.shtml>
5. IPAMS. *Invasive Plant Atlas of the MidSouth*. Available at: <http://gri.msstate.edu/ipams>
6. IPANE. *Invasive Plant Atlas of New Engla* Available at: <http://nbii-nin.ciesin.columbia.edu/ipane/>
7. Jepson eFlora. *Rosa rubiginosa, Jepson Flora Project*. Available at: <http://ucjeps.berkeley.edu/IJM.html>
8. NCU Flora. *Flora of the Southeastern United States*. Available at: <http://herbarium.unc.edu/seflora/firstviewer.htm>
9. NMBCC. *New Mexico Biodiversity Collections Consortium Gateway to New Mexico Biodiversity*. Available at: <http://NMBiodiversity.org>
10. PNW Herbaria. *Consortium of Pacific Northwest Herbaria*. Available at: <http://www.pnwherbaria.org>
11. SWEMP. *Southwest Exotic Mapping Program*. Available at: <http://sbsc.wr.usgs.gov/research/projects/swepic/swemp/swempA.asp>

### Europe

12. Anthos. *Information System of Plants in Spain*. Available at: <http://www.anthos.es>
13. Artamonov AA. New records of *Rosa rubiginosa* L. in central Russia. *Byull Mosk Obs Ispyt Prir Biol*. 2002; 107: 54–6.
14. Artportalen. *Reporting System for Vascular Plants and Fungi*. Available at: <http://www.artportalen.se/default.asp>
15. Artsdatabanken. *Nasjonal kunnskapskilde for biologisk mangfold*. Available at: <http://www.artsdatabanken.no/>

16. Benkert D, Fukarek F, Korsch H. Verbreitungsatlas der Farn- und Blütenpflanzen Ostdeutschlands. Jena: Fischer; 1996.
17. Bolós O, Vigo J. Flora dels Països Catalans. Barcelona: Editorial Barcino; 1995.
18. Dubovik ON. Hrupy Rosa tomentosa Smith i R. rubiginosa L. u flori Ukrayiny. Ukr Bot Zhurnal. 1989;46:16–20.
19. FCD. *Flora Croatica Database*. Available at: <http://hirc.botanic.hr/fcd/>
20. Flora Italiana. *Flora Italiana*. Available at: <http://www.homolaicus.com/scienza/erbario/utility/floraitalica/flora/index.htm>
21. Geydeman TS. Opređelitel' vysshikh rasteniy Moldavskoy SSR. Kishinev: Stiinta; 1986.
22. Gubanov IA, Kiseleva K V, Novikov VS, Tikhomirov VN. Illustrated Manual of the Middle Russia Plants. Moscow: KMK; 2003.
23. Haeupler H, Jagel A, Schumacher W. Verbreitungsatlas der Farn- und Blütenpflanzen in Nordrhein- Westfalen. 616: Landesanstalt für Ökologie, Bodenordnung und Forsten; 2003.
24. Haeupler H, Schönfelder P. Atlas der Farn- und Blütenpflanzen der Bundesrepublik Deutschland. Stuttgart : Ulmer; 1988.
25. Hultén E. Atlas of the distribution of vascular plants in northwestern Europe. Stockholm: Generalstabens Litografiska Anstalts Förlag; 1971.
26. ICEL. *Maps of Icelandic Plants*. Available at: <http://vefsja.ni.is/website/plontuvefsja>
27. INPN. *Inventaire National du Patrimoine Naturel*. Available at: <http://inpn.mnhn.fr/accueil/index?lg=en>
28. Kasviatlas. *Atlas of the Vascular Flora of Finla* Available at: <http://www.luomus.fi/kasviatlas>
29. Lang W, Wolff P. Flora der Pfalz. Verbreitungsatlas der Farn- und Blütenpflanzen für die Pfalz und ihre Randgebiete. Speyer: Veröffentlichung Pfälzische Gesellschaft zur Förderung der Wissenschaften; 1993.
30. Maevskii PF. Flora of middle belt of the European part of the USSR. Moscow: Selkhozgiz; 1954.
31. Majorov SR, Kramina TE. Notes on distribution of some Rosaceae Juss. in eastern Europe. Byull Mosk Obs Ispyt Prir Biol. 2003;108:50–4.
32. 2. Meusel H, Jäger EJ, Weinert E. Vergleichende Chorologie der zentraleuropäischen Flora. Jena: Fischer; 1965.
33. NBN Gateway. *National Biodiversity Network's Gateway*. Available at: <http://www.searchnbn.net/>
34. Poldini L. Atlante corologico delle piante vascolari nel Friuli-Venezia Giulia. Udine: Direzione Regionale delle Foreste e dei Parchi; 1991.



35. SILENE. *Système d'Information et de Localisation des Espèces Natives et Envahissantes*. Available at: <http://silene.eu/index.php?cont=accueil>
36. Raabe EW. *Atlas der Flora Schleswig-Holsteins und Hamburgs*. Neumünster: Wachholtz; 1987.
37. Sauer E. *Die Gefäßpflanzen des Saarlandes*. Saarbrücken: Aus Natur und Landschaft; 1993.
38. Schumacher W. *Atlas der Farn- und Blütenpflanzen des Rheinlandes*. Bonn: Landwirtschaftliche Fak. der Rheinischen Friedrich-Wilhelms-Univ.; 1995.
39. Schönfelder P, Bresinsky A. *Verbreitungsatlas der Farn- und Blütenpflanzen Bayerns*. Stuttgart: Ulmer; 1990.
40. Slavík B, Smejkal M, Dvořáková M, Grulich V. *Květena České republiky*. Praha; 1995.
41. Sokolov SI, Svjaseva OA, Kubli VA. *Ranges of trees and shrubs of the USSR*. Leningrad: Nauka; 1980.
42. SOPHY. *Banque de Données Botaniques et Ecologiques*. Available at: <http://sophy.u-3mrs.fr/sophy.htm>
43. Tzvelev NN. *Keys to the vascular plants of North-West Russia (Leningrad, Pskov and Novgorod Regions)*. St. Petersburg: SPbSCPA press; 2000.
44. Tutin TG, Heywood VH, Burges NA, Moore DM, Walters SM, Webb DA. *Flora Europaea*, Vol. 4, Plantaginaceae to Compositae (and Rubiaceae). Cambridge: Cambridge University Press; 1976.
45. Tzvelev NN. *Flora of East Europe*. St. Petersburg; 2001.
46. Yuzepchuk S V. Rose (wild rose) - *Rosa* L. In: Shishkin BK, Yuzepchuk S V, editors. *Flora URSS*. Moscow, Leningrad; 1941. p. 431–506.
47. Zielinski J. *Rosa rubiginosa* L. and *R. heckeliana* Tratt. in Bulgaria. *Fragm Florist Geobot.* 1986;30:207–11.
48. Zinovskis RE, Bize MA, Knappe DA, Kucheneva GG. *Conspectus of dendroflora of Kaliningrad region*. Zinatne; 1983.

## Asia

49. Galuško AI. *Flora Severnogo Kavkaza*. Rostow: Isd. Rostovskogo Univ.; 1980.
50. Grossgejm AA. *Opredelitel' rastenij Kavkaza*. Moskow: Sovj. Nauk; 1949.

## Australia and New Zealand

51. AVH. *Australia's Virtual Herbarium*. Available at: <http://www.chah.gov.au/avh/>
52. Allan Herbarium (2000) *Ngā Tipu o Aotearoa - New Zealand Plant Names Database*. Available at: <http://nzflora.landcareresearch.co.nz>

## South America

53. Correa MN. Flora Patagónica. Buenos Aires: Colección Científica del INTA
54. Damascos MA, Bran D. Morfología, distribución y abundancia de tres especies de rosas invasoras en la región Andino-Patagónica de Argentina. San Carlos de Bariloche: Resúmenes: Jornadas Argentinas de Botánica; 2003.
55. Damascos MA, Gallopin GG. Ecología de un arbusto introducido (*Rosa rubiginosa* L. = *Rosa eglanteria* L.): riesgo de invasión y efectos en las comunidades vegetales de la región andino-patagónica de Argentina. Rev Chil Hist Nat. 1992; 65: 395–407.
56. Zuloaga FO, Morrone O. Catálogo de las plantas vasculares de la República Argentina. St. Louis: Missouri Botanical Garden Press; 1999.
57. Quiroz C, Pauchard A, Marticorena A, Cavieres LA. Manual de plantas invasoras del centro-sur de Chile. Laboratorio de Invasiones Biológicas; 2009.

## Africa

58. AGIS. *Agricultural Geo-Referenced Information System*. Available at: <http://www.agis.agric.za>
59. APD. *African Plants Database*. Available at: <http://www.ville-ge.ch/musinfo/bd/cjb/africa/recherche.php?langue=an>
60. Buzunova I, Romo A. *Rosa rubiginosa* (Rosaceae) in Morocco-first records from northern Africa. Dendrobiology. 2011;66:99–103.
61. SANBI. *South African National Biodiversity Institute*. Available at: <http://www.sanbi.org/>